



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.



Published by the contributors to advance the Science of cold-blooded vertebrates.

FISHES AND REPTILES FROM ASSUAN, EGYPT.

Mr. C. F. Silvester recently submitted the following collection :

FISHES.

- Anguilla anguilla* (Linnæus). Adult.
Hydrocyon forskalii Cuvier. Adult and half-grown.
Alestes baremose (Joannis). One adult.
Labeo coubie Rüppell. Adult.
Barbus bynni (Forskål). Small example.
Eutropius niloticus (Rüppell). Adult.
Siluranadon auritus (G. St. Hilaire). Two adults.
Porcus bajad (Forskål). Adult.
Synodontis schall (Schneider). One adult.
Clarias lazera Valenciennes. Adult and young.
Lates niloticus (Linnæus). Adult.
Tilapia nilotica (Linnæus). Adult.

REPTILES.

- Eryx thebaicus* Reuss. Four.
Eryx jaculus (Linnæus). Eight.
Coluber diadema Schlegel. Two adults.
Coluber florulentis (G. St. Hilaire). Eight, all ages.
Psammophis sibilans (Linnæus). Seven, adult and young.
Psammophis shokari (Forskål). Six, adult and young.
Naja haje (Linnæus). Four adults.

Cerastes cerastes (Linnæus). Two adults.
Ptyodactylus hasselquistii (Donndorff). One adult.
Tarentola annularis (G. St. Hilaire). Four adults.
Agama flavimaculata (Rüppell). Adult.
Agama pallida Reuss. Three adults.
Uromastix ornatus Rüppell. One adult. This species
 apparently not previously recorded definitely
 from Egypt. Back uniform in color.
Varanus griseus (Daudin). Adult.
Varanus niloticus (Linnæus). Adult.
Acanthodactylus boskianus (Daudin). One.
Scincus scincus (Linnæus). Six, mostly adults.

HENRY W. FOWLER,
Philadelphia, Pa.

AMBLYSTOMA OF LONG ISLAND

Three species of *Amblystoma* (*A. punctatum*,
A. opacum, *A. tigrinum*) have been listed as being
 found in the vicinity of New York City.* Of these
 only *A. tigrinum* is mentioned, without specific
 data, as occurring on Long Island. Observations
 by the writer and material in the Museum of the
 Brooklyn Institute of Arts and Sciences show the
 following Long Island records:

AMBLYSTOMA PUNCTATUM (Linnæus).

(Spotted Salamander).

2 specimens Cold Spring Harbor, L. I. Aug., 1900.
 —G. K. Cherrie and L. Ericson.

3 specimens Forest Park, Brooklyn. May, 1906.
 Under stones and logs, damp soil.
 —Geo. P. Engelhardt.

1 specimen Huntington, L. I. Aug., 1913.
 In potato patch. —Miss Graeve.

*Sherwood, William L. Salamanders found in the vicinity of New York City.
 Linn. Soc. 1894-95. No. 7.

Ditmars, Raymond L. The Batrachians of the vicinity of New York City.
 Amer. Mus. Journal, Vol. V, No. 4. Oct., 1905.